Rick D. Russotto, Thomas P. Ackerman, and Dale R. Durran Department of Atmospheric Sciences, University of Washington, Seattle, WA, USA russotto@atmos.washington.edu, durrand@atmos.washington.edu, ackerman@atmos.washington.edu

## I. OBJECTIVES

Existing cloud-resolving models of thin cirrus in the tropical tropopause layer unrealistically assumed all ice crystals were spherical. We want to know how sensitive these clouds are to ice crystal shape, and to better understand the physics of their life cycle.

## II. METHODS

Model:	System for Atmospheric Modeling (SAM)		
<b>Resolution:</b>	Δx = 100 m, Δz = 25 m, Δt = 6 s		
Initial ice crystal size:	3.0 µm radius, or equivalent mass for non-spheres		
Bin micro- physics scheme:	Dinh & Durran [2012], modified for oblate and prolate sphe- roids with aspect ratios of 6		
Radiation scheme:	Plane-parallel radiation solver in SAM; absorption coefficients for spheres, plates and columns parameterized using RRTM radiative transfer model		

### **Table 1:** Run Names and Assumptions

Run Name	Fall Speed	Growth Rate	LW Absorption	SW Absorption
Control	Spheres	Spheres	Spheres	Spheres
Oblate Fall Speed	Oblate	Spheres	Spheres	Spheres
Oblate Growth Rate	Spheres	Oblate	Spheres	Spheres
Oblate Microphysics	Oblate	Oblate	Spheres	Spheres
Oblate-Full	Oblate	Oblate	Plates	Spheres
Zero Fall Speed	0	Spheres	Spheres	Spheres
Double LW Absorption	Spheres	Spheres	Spheres x 2.0	Spheres
Half LW Absorption	Spheres	Spheres	Spheres x 0.5	Spheres
Zero SW Absorption	Spheres	Spheres	0	Spheres
Zero Radiative Absorption	Spheres	Spheres	0	0

Results from prolate spheroid runs not included because of similarity to oblate spheroids.

# Sensitivity of Thin Tropical Cirrus Clouds to Ice Crystal Shape and Radiative Absorption

III. RESULTS







## V. CONCLUSIONS

Thin tropical tropopause layer cirrus are sensitive to the amount of LW radiation absorbed, which implies a dependence on underlying clouds, humidity and surface temperature. Sensitivity to ice crystal shape is minimal.





### IV. DISCUSSION

• Time evolution of the cloud (Figure 1) is qualitatively consistent with previous work [Dinh et al., 2010] Radiatively induced circulation deforms cloud and maintains it against sedimentation • Small scale convection develops at the cloud top

• Shape has little effect on center of cloud ice mass (Figure 2A) or total ice mass (Figure 2B)

• Spheroids have 1/3 lower fall speeds than spheres • Centers of mass in Control, Oblate Fall Speed and Zero Fall Speed cases are consistent with this (Figure 3)

• Stronger radiative absorption versus Control leads to • Higher center of mass (Figure 4A) • A *less* massive cloud (Figure 4B), due to • Lower relative humidity in cloud and source air Entrainment of dry air by more vigorous convection

#### ACKNOWLEDGEMENTS

This research was supported by the National Science Foundation (NSF) through grant ATM-0926996 and by the U.S. Department of Defense (DoD) through the National Defense Science and Engineering Graduate Fellowship (NDSEG) Program. Model simulations were performed using the Hyak supercomputer system at the University of Washington. The authors are grateful to Tra Dinh, who provided valuable insights based on her own experience modeling TTL cirrus, and to Stephen Warren, who provided helpful sugges tions for calculating the radiative properties of non-spherical ice crystals.

#### REFERENCES

Dinh, T., and D.R. Durran (2012), A hybrid bin scheme to solve the condensation/evaporation equation using a cubic distribution function, Atmospheric Chemistry and Physics, 12 (2), 1003-1011, doi:10.5194/acp-12-1003-2012.

Dinh, T., D. Durran, and T. Ackerman (2010), Maintenance of tropical tropopause layer cirrus, Journal of Geophysical Research (Atmospheres), 115, D02104, doi:10.1029/2009JD012735

Russotto, R.D., T.P. Ackerman, and D.R. Durran (in preparation), Sensitivity of Thin Tropical Cirrus Clouds to Ice Crystal Shape and Radiative Absorption, Journal of Geophysical Research (Atmospheres).